

## ORIGINAL ARTICLE

# The *Arabidopsis thaliana* gene *AtERF019* negatively regulates plant resistance to *Phytophthora parasitica* by suppressing PAMP-triggered immunity

Wenqin Lu<sup>1</sup> | Fengyan Deng<sup>2</sup> | Jinbu Jia<sup>1,3</sup> | Xiaokang Chen<sup>4</sup> | Jinfang Li<sup>1</sup> |  
 Qujiang Wen<sup>1</sup> | Tingting Li<sup>5</sup> | Yuling Meng<sup>5</sup> | Weixing Shan <sup>5</sup>

<sup>1</sup>State Key Laboratory of Crop Stress Biology for Arid Areas and College of Plant Protection, Northwest A&F University, Yangling, China

<sup>2</sup>State Key Laboratory of Crop Stress Biology for Arid Areas and College of Life Sciences, Northwest A&F University, Yangling, China

<sup>3</sup>Institute of Plant and Food Science, Department of Biology, Southern University of Science and Technology, Shenzhen, China

<sup>4</sup>State Key Laboratory of Crop Stress Biology for Arid Areas and College of Horticulture, Northwest A&F University, Yangling, China

<sup>5</sup>State Key Laboratory of Crop Stress Biology for Arid Areas and College of Agronomy, Northwest A&F University, Yangling, China

## Correspondence

Weixing Shan and Yuling Meng, State Key Laboratory of Crop Stress Biology for Arid Areas and College of Agronomy, Northwest A&F University, Yangling, Shaanxi 712100, China.

Emails: wxshan@nwfau.edu.cn; mengyuling@nwfau.edu.cn

## Funding information

National Natural Science Foundation of China, Grant/Award Number: 31125022 and 31930094; China Agricultural Research System, Grant/Award Number: CARS-09; Programme of Introducing Talents of Innovative Discipline to Universities (project 111) from the State Administration of Foreign Experts Affairs, Grant/Award Number: B18042

## Abstract

*Phytophthora* species are destructive plant pathogens that cause significant crop losses worldwide. To understand plant susceptibility to oomycete pathogens and to explore novel disease resistance strategies, we employed the *Arabidopsis thaliana*-*Phytophthora parasitica* model pathosystem and screened for *A. thaliana* T-DNA insertion mutant lines resistant to *P. parasitica*. This led to the identification of the resistant mutant 267-31, which carries two T-DNA insertion sites in the promoter region of the *ethylene-responsive factor 19* gene (*ERF019*). Quantitative reverse transcription PCR (RT-qPCR) assays showed that the expression of *ERF019* was induced during *P. parasitica* infection in the wild type, which was suppressed in the 267-31 mutant. Additional *erf019* mutants were generated using CRISPR/Cas9 technology and were confirmed to have increased resistance to *P. parasitica*. In contrast, *ERF019* overexpression lines were more susceptible. Transient overexpression assays in *Nicotiana benthamiana* showed that the nuclear localization of *ERF019* is crucial for its susceptible function. RT-qPCR analyses showed that the expression of marker genes for multiple defence pathways was significantly up-regulated in the mutant compared with the wild type during infection. Flg22-induced hydrogen peroxide accumulation and reactive oxygen species burst were impaired in *ERF019* overexpression lines, and flg22-induced MAPK activation was enhanced in *erf019* mutants. Moreover, transient overexpression of *ERF019* strongly suppressed INF-triggered cell death in *N. benthamiana*. These results reveal the importance of *ERF019* in mediating plant susceptibility to *P. parasitica* through suppression of pathogen-associated molecular pattern-triggered immunity.

## KEYWORDS

*Arabidopsis thaliana*, oomycete, *Phytophthora parasitica*, susceptibility, transcription factor

Wenqin Lu and Fengyan Deng contributed equally.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Molecular Plant Pathology* published by British Society for Plant Pathology and John Wiley & Sons Ltd

## 1 | INTRODUCTION

The “plant destroyer” *Phytophthora* causes devastating disease in a large number of crops and forest seedlings worldwide. For example, potato late blight caused by *Phytophthora infestans* can lead to severe decreases in production and also serious economic losses (Haverkort *et al.*, 2008). *P. sojae*, *P. ramorum*, *P. parasitica*, and *P. capsici* can also cause important agricultural diseases such as soybean root rot and oak stagnation (Tyler, 2002; Grünwald *et al.*, 2012; Lamour *et al.*, 2012; Meng *et al.*, 2014; Kamoun *et al.*, 2015; Panabières *et al.*, 2016).

Plants have sufficient weapons to repel pathogen attacks, but need to recognize the pathogen in time, which mainly occurs through two different systems. One is referred to as PAMP-triggered immunity (PTI), which is activated by transmembrane pattern-recognition receptors (PRRs) through recognition of pathogen- or microbe-associated molecular patterns (PAMPs or MAMPs) (Jones and Dangl, 2006; Boller and Felix, 2009; Bigeard *et al.*, 2015; Boutrot and Zipfel, 2017) and initiates a series of immune responses, including reactive oxygen species (ROS) bursts, callose deposition, biosynthesis of phytohormones (such as salicylic acid [SA], jasmonate [JA], and ethylene [ET]), and the expression of a large number of defence-related genes (Zipfel *et al.*, 2004; Naito *et al.*, 2008). The other system is called effector-triggered immunity (ETI), which is based on the specific recognition between pathogen effectors and plant resistance (R) proteins, according to the gene-for-gene theory. This recognition leads to a rapid and localized hypersensitive response (HR, cell death) at infection sites and inhibits pathogen colonization (Jones and Dangl, 2006).

Although priming of immunity responses when pathogens attack is the key to resistance, excessive and inappropriate defence responses interfere with the growth of a plant. To mitigate the trade-off between growth and defence, plants have evolved a series of mechanisms to negatively regulate defence pathways. For example, the rice *Pigm* locus confers durable resistance to the fungus *Magnaporthe oryzae* without yield penalty; this is achieved through epigenetic regulation of two antagonistic receptors, *PigmR* and *PigmS*, encoded by this locus (Deng *et al.*, 2017). *PigmR* confers broad-spectrum resistance, whereas *PigmS* competitively attenuates *PigmR* homodimerization to suppress resistance (Deng *et al.*, 2017). In addition, growth-related hormones, auxin, brassinosteroids (BRs), and gibberellins (GAs) can directly or indirectly negatively regulate PTI-mediated defence (Yamada, 1993; Chen *et al.*, 2007; Albrecht *et al.*, 2012; Jaillais and Vert, 2012). However, these negative regulatory pathways can be hijacked by effectors secreted by pathogens to promote infection (Jones and Dangl, 2006). For example, plant cinnamyl alcohol dehydrogenase 7 (CAD7), which is involved in the negative regulation of plant resistance to *Phytophthora* pathogens including *P. infestans*, *P. paratisica*, and *P. capsici*, is a common target of multiple AVR3a-like effectors from *Phytophthora* pathogens. These effectors suppress PTI responses by stabilizing CAD7 (Li *et al.*, 2019). In potato, StVIK and StKRBP1 are targeted by the RXLR effectors Pi17316 and Pi04089 from *P. infestans*, respectively, to facilitate invasion (Wang *et al.*, 2015; Murphy *et al.*, 2018). Therefore,

appropriate manipulation of negative regulators of plant immunity has the potential to improve broad-spectrum disease resistance.

To explore the mechanisms by which negative regulators suppress plant resistance to pathogens, we used a model compatible system between *Arabidopsis thaliana* and the oomycete pathogen *P. parasitica* (Wang *et al.*, 2011b) to screen for *A. thaliana* T-DNA insertion mutants resistant to *P. parasitica* infection. We identified an *erf019* mutant that showed less susceptibility to *P. parasitica*. Our analysis revealed that *ERF019* negatively regulates plant defence responses to *Phytophthora* pathogens by suppressing PAMP-triggered immunity, thus acting as an important regulator in balancing plant disease resistance and growth.

## 2 | RESULTS


### 2.1 | Identification of an *erf019* mutant, 267-31, that limits colonization of *P. parasitica*

To identify genes that negatively regulate defence against *P. parasitica* infection, we screened nearly 10,000 independent *Arabidopsis* T-DNA insertion lines (Zhang *et al.*, 2005) to identify mutants involved in limiting the colonization of *P. parasitica*. This led to the identification of the mutant 267-31 (Figure 1a,b). In comparison to wild-type Col-0, growth of the pathogen, *P. parasitica* Pp016 (Wang *et al.*, 2011b; Zhang *et al.*, 2011), was much more restricted in 267-31 at 3 days postinoculation (dpi) (Figure 1a). Consistent with this finding, quantification of *P. parasitica* colonization in infected *Arabidopsis* leaves revealed less colonization in 267-31 (Figure 1b).

Thermal asymmetric interlaced (TAIL)-PCR (Liu *et al.*, 1995) was applied to obtain sequences flanking the T-DNA insertion sites in 267-31. Sequence analysis showed that there were two T-DNA insertion sites located 309 and 376 bp upstream of the *ERF019* translation start codon, respectively (Figure 1c). The two T-DNA insertion fragments were adjacent and in opposite orientations. Quantitative reverse transcription (RT) PCR (RT-qPCR) analysis showed that the *ERF019* transcript levels were dramatically lower in 267-31 than in the wild-type Col-0 (Figure 1d). Thus, *ERF019* might play a negative role in resistance against *P. parasitica*.

### 2.2 | *ERF019* contributes to plant susceptibility to *P. parasitica*


To confirm that *ERF019* contributes to plant susceptibility to *P. parasitica*, we used the CRISPR/Cas9-mediated genome-editing tool to knock out the *ERF019* gene. Two target sites in the exon of *ERF019* were chosen (Figure 1e), and the corresponding sgRNA/Cas9 vectors were transformed into the wild-type Col-0 via *Agrobacterium tumefaciens*-mediated transformation. The mutations at the target sites in the CRISPR/Cas9 transformants were examined using PCR and DNA sequencing analysis, which showed that approximately 90% of T<sub>0</sub> transformants carried mutations at one sgRNA target site



**FIGURE 1** The *erf019* mutants of *Arabidopsis thaliana* limit *Phytophthora parasitica* colonization. (a) Trypan blue staining showing the disease symptoms of the *erf019* T-DNA insertion mutant 267-31 and the wild-type Col-0 infected with *P. parasitica* strain Pp016. The concentration of zoospore suspensions was adjusted to 200 zoospores/ $\mu$ l. Detached leaves of 4-week-old *A. thaliana* seedlings were drop-inoculated with 10  $\mu$ l *P. parasitica* zoospores (200 zoospores/ $\mu$ l) and photographed at 3 days postinoculation (dpi). (b) Quantitative reverse transcription PCR (RT-qPCR) quantification of pathogen colonization. Total genomic DNA from *P. parasitica*-infected regions was isolated at 2 dpi. Quantitative PCR (qPCR) with primers specific for the *A. thaliana* *UBC9* gene (*AtUBC*) and the *P. parasitica* *UBC* gene (*PpUBC*) was used to determine *P. parasitica* biomass in infected plant tissues. The relative *P. parasitica* biomass was calculated by *PpUBC/AtUBC* and normalized using the value of Col-0. (c) Two T-DNAs were inserted in the promoter region of *ERF019*. (d) RT-qPCR analysis to quantify the expression of *ERF019* in 2-week-old seedlings of *Arabidopsis* mutant 267-31 and wild-type Col-0. *UBC9* was used as the internal control. Bars represent standard errors from three biological replicates and asterisks indicate statistical significance based on *t* test (\*\**p* < .01). (e) Targeted indel mutations at the *ERF019* gene. Representative sequences of CRISPR/Cas9-based knockout mutant alleles identified from transgenic plants expressing sgRNA targeting *ERF019*. (f) Disease symptoms of CRISPR/Cas9-based knockout mutants and the wild-type Col-0 infected with *P. parasitica*. The concentration of zoospore suspensions was adjusted to 200 zoospores/ $\mu$ l. Detached leaves of 4-week-old *Arabidopsis* plants were drop-inoculated with 20  $\mu$ l *P. parasitica* zoospores (200 zoospores/ $\mu$ l) and photographed at 2 dpi. (g) Pathogen colonization in knockout lines. Detached leaves of 4-week-old *Arabidopsis* were drop-inoculated with 20  $\mu$ l zoospores of *P. parasitica* transformant 1121, which stably expresses green fluorescent protein (GFP), and visualized under a fluorescence microscope at 2 dpi. Green fluorescence indicates *P. parasitica* hyphae, autofluorescence from leaf tissue is visible as red signal. The white bars indicate 500  $\mu$ m. (h) Quantification of *P. parasitica* biomass in inoculated leaves of knockout lines by qPCR. Error bars represent SD, and asterisks indicate statistical significance based on *t* test (\**p* < .05; \*\**p* < .01). Similar results were obtained from at least three individual experiments

while no mutations were found at the other target site. Of three individual homozygous knockout lines chosen for further analysis, two contain a 1-bp insertion (ko13-4 and ko58-7) and one contains a 4-bp deletion (ko26-14) in the coding region of *ERF019* (Figure 1e), which results in a frameshift mutation and predicted truncated protein (Figure S1). Phenotypic observations revealed that none of these three *erf019* knockout lines had obvious morphological abnormalities (Figure S2), suggesting *ERF019* is not essential for plant growth and development. When inoculated with 1121, a transformant of Pp016 stably and widely expressing ER-rendered green fluorescent protein (GFP) under control of the constitutive *Hsp70* promoter of *Bremia lactucae* (Zhang *et al.*, 2011), the CRISPR/Cas9-edited mutants had attenuated *P. parasitica* leaf colonization, with significantly smaller water-soaked lesions compared with those in the wild-type Col-0 (Figure 1f). Microscopic observation showed that fewer GFP-expressing hyphae colonized the CRISPR/Cas9-edited mutants compared with the wild-type Col-0 (Figure 1g). In addition, quantification of *P. parasitica* biomass revealed that CRISPR/Cas9-edited mutants exhibited limited colonization of *P. parasitica* (Figure 1h).

To further clarify the function of *ERF019* in plant susceptibility, we transformed an *ERF019*-overexpression (OE) construct, in which *ERF019* expression is under the control of constitutive cauliflower mosaic virus (CaMV) 35S promoter, into the Col-0 background. Three *ERF019*-OE lines, OE71, OE72, and OE74, were selected for further analysis following confirmation by RT-qPCR analysis that the *ERF019* transcript levels were significantly increased in these lines (Figure 2d). Infection assays with *P. parasitica* revealed that *ERF019*-OE plants were more susceptible than the wild-type Col-0. Two days after pathogen infection, *ERF019*-OE lines developed much larger water-soaked lesions than the wild-type Col-0 (Figure 2a). Heavier hyphal colonization was also visible in *ERF019*-OE plants when infected with the *P. parasitica* transformant 1121, which stably expresses GFP (Figure 2b). In addition, both a trypan blue staining assay and a quantification of *P. parasitica* biomass revealed that *ERF019*-OE plants exhibited enhanced disease susceptibility to *P. parasitica* (Figure 2c,e). These results confirmed that *ERF019* negatively regulates plant resistance to *P. parasitica*.



**FIGURE 2** The *ERF019*-overexpression (*ERF019*-OE) *Arabidopsis thaliana* plants were more susceptible to *Phytophthora parasitica*. (a) Disease symptoms of *ERF019*-OE plants infected with 10  $\mu$ l *P. parasitica* zoospores (200 zoospores/ $\mu$ l) and photographed at 2 days postinoculation (dpi). The concentration of zoospore suspensions was adjusted to 200 zoospores/ $\mu$ l using microscopy. (b) Pathogen colonization on *ERF019*-OE plants. Detached leaves of 4-week-old *A. thaliana* plants were drop-inoculated with 10  $\mu$ l zoospores of *P. parasitica* transformant 1121, which stably expresses green fluorescent protein (GFP), and visualized under a fluorescence microscope at 2 dpi. Green fluorescence indicates *Phytophthora* hyphae. (c) Trypan blue staining of *ERF019*-OE plants infected with *P. parasitica*. Plant cells infected by pathogen were coloured. (D) Quantitative reverse transcription PCR analysis showed the accumulation of *ERF019* transcripts in rosette leaves of *ERF019*-OE plants. Data represent the ratio of *ERF019* expression between *ERF019*-OE plants and wild-type Col-0. *UBC9* was used as the internal control. Bars represent standard errors from three biological replicates and asterisks indicate statistical significance based on t test (\*\* $p < .01$ ). (e) Quantification of *P. parasitica* biomass in inoculated leaves of *ERF019*-OE plants by quantitative PCR. Error bars represent SD, and asterisks indicate statistical significance based on t test (\* $p < .05$ ; \*\* $p < .01$ ). Similar results were obtained from at least three individual experiments

### 2.3 | *ERF019* expression is induced on *P. parasitica* infection

To examine whether the expression of *ERF019* is responsive to *P. parasitica* infection, we used RT-qPCR to measure the *ERF019* expression levels in *P. parasitica*-inoculated leaves at different time points. The results showed that in Col-0, *ERF019* expression was highly induced at 3 hours postinoculation (hpi) and slightly induced at 6 and 12 hpi compared with that in the uninfected leaves and the mock-inoculated controls. The expression levels of *ERF019* at 24 hpi were decreased to a level similar to that of the uninfected leaves, though it appeared a little higher than that in the mock-inoculated controls. These observations indicate that *ERF019* is responsive to *P. parasitica* infection. In contrast, the expression level of *ERF019* in the 267-31 mutant was significantly lower than that in the wild type, both in the uninfected and infected leaves, though the mutant still produced few detectable transcripts (Figure 3).

### 2.4 | Nuclear localization of *ERF019* is required for its susceptibility function


*ERF019* contains an APETALA2 (AP2) domain and is, therefore, predicted as a member of the ethylene-responsive factor (ERF)/AP2 transcription factor family. We transiently co-expressed the *ERF019*-GFP fusion construct with the nucleus marker H2B-mCherry in *Nicotiana benthamiana* leaves and found that *ERF019*-GFP can be observed in both nucleus and cytoplasm (Figure S3). To confirm that the nuclear localization of *ERF019* is necessary for its function in promoting pathogen infection, we altered the subcellular localization of *ERF019* by fusing it with a nuclear export sequence (NES). A construct expressing *ERF019* fused with a mutant NES (nes) was used as a control. We first transiently expressed the *ERF019*-GFP-NES and *ERF019*-GFP-nes chimeric proteins under the control of the CaMV 35S promoter in *N. benthamiana* leaves and observed the subcellular localization by fluorescence microscopy. *ERF019*-GFP-NES was clearly exported to the cytosol and was barely detectable in the nucleus, while *ERF019*-GFP-nes displayed a localization pattern similar


to that of *ERF019*-GFP (Figure S4), which indicates that the NES was functional. Next, to examine whether the nuclear export of *ERF019* affects its role in plant susceptibility, we transiently expressed these fusion proteins in *N. benthamiana* leaves and challenged the leaves with *P. parasitica*. Notably, leaves transiently expressing *ERF019*-GFP displayed significantly larger infection lesions compared with those expressing the FLAG-GFP control (Figure 4), which is consistent with the role of *ERF019* as a negative regulator of plant immunity. *ERF019*-GFP-NES, but not *ERF019*-GFP-nes, lost its ability to enhance *P. parasitica* colonization of *N. benthamiana* (Figure 4). These results suggest that the nuclear localization of *ERF019* is required for its function in promoting colonization of *P. parasitica*.

### 2.5 | Expression of defence marker genes is up-regulated in the *erf019* mutant

Because *ERF019* negatively regulates plant resistance to *P. parasitica*, we examined the potential role of *ERF019* in known defence pathways by measuring the expression of marker genes associated with these pathways in 267-31 and the CRISPR/Cas9-edited *erf019* mutant lines (ko13-4, ko26-14). *ICS1* (*Isochorismate synthase1*) and *PAL1* (*Phenylalanine ammonia lyase 1*) are involved in the synthesis of SA (Wildermuth *et al.*, 2001), and *PR1* (*Pathogenesis-related gene 1*) is a well-established marker gene for the SA signalling pathway (Uknes *et al.*, 1992). *LOX2* (*Lipoxygenase2*) is involved in the synthesis of jasmonic acid (JA) (Sasaki *et al.*, 2001), and *PDF1.2* (*Plant defensin gene 1.2*) and *VSP2* (*Vegetative storage protein 2*) are genes that respond to JA (Pieterse *et al.*, 2009). *FRK1* (*Flg22-induced receptor-like kinase 1*) is a core gene that is induced by a conserved 22 amino acid epitope from bacterial flagellin (flg22), and is frequently used to monitor PTI (Shan *et al.*, 2008). *ACS2* and *ACS6*, two members of the ACS gene family that encode 1-amino-cyclopropane-1-carboxylase synthase, are involved in the ethylene biosynthesis (Van der Straeten *et al.*, 1992), and *EIN2* (*Ethylene-insensitive protein 2*) is an important regulator in the ET signalling pathway (Alonso *et al.*, 1999) and *ERF6* (*Ethylene-responsive factor 6*) is an ET-related signalling gene (Moffat *et al.*, 2012). An RT-qPCR assay showed that the expression of defence-related genes was altered in

**FIGURE 3** Quantitative reverse transcription PCR analysis for the expression of *ERF019* in *Arabidopsis thaliana* Col-0 and the T-DNA insertion mutant 267-31 at different time points after *Phytophthora parasitica* inoculation. *AtUBC9* was used as the internal control. Bars represent standard errors from three biological replicates and asterisks indicate statistical significance based on *t* test (\* $p < .05$ ; \*\* $p < .01$ ; ns, not significant). 0 h, uninfected leaves; hpi, hours postinoculation






**FIGURE 4** Nuclear localization is required for ERF019 to promote *Phytophthora parasitica* growth in *Nicotiana benthamiana*. *P. parasitica* colonization of *N. benthamiana* leaves expressing FLAG-GFP, ERF019-GFP, ERF019-GFP-NES or ERF019-GFP-nes. *Agrobacterium tumefaciens* GV3101 cells carrying FLAG-GFP, ERF019-GFP, ERF019-GFP-NES or ERF019-GFP-nes were infiltrated into leaves of *N. benthamiana*, and infiltrated leaves were challenged with 10  $\mu$ l *P. parasitica* Pp016 zoospores (200 zoospores/ $\mu$ l) 48 hr after infiltration. The concentration of zoospore suspension was adjusted to 200 zoospores/ $\mu$ l. Disease symptoms were observed 2 days postinoculation and leaves were stained with trypan blue. At least 10 leaves were used for the test. Error bars represent SD, and asterisks indicate statistical significance based on a two-tailed t test (\* $p < .05$ ; \*\* $p < .01$ ). Similar results were obtained for at least three individual experiments

the *erf019* mutants compared with the wild-type Col-0. In *P. parasitica*-inoculated plants, the expression levels of *ICS1*, *PR1*, *VSP2*, *LOX2*, *PDF1.2*, and *FRK1* in 267-31 and the CRISPR/Cas9-edited lines were significantly higher than that in the wild-type Col-0 (Figure 5). The expression of *PAL1* was down-regulated on infection, and showed similar level between mutants and the wild-type Col-0 (Figure 5). However, for the marker genes in the ET signalling pathway, the expression levels of *ACS6*, *EIN2*, and *ERF6* appeared similar between mutants and the wild type, although *ACS2* was induced to higher levels in mutants at some time points (Figure 5). Taken together, these results indicate that *ERF019* may play an important role in the SA and JA defence signalling pathways but not the ET signalling pathway.

## 2.6 | ERF019 suppresses PTI responses

Perception of flg22 triggers a series of immunity responses, including an oxidative burst, rapid and transient accumulation of ROS, and

the activation of MAP kinases. *ERF019* was reported to be highly induced by flg22 (Sano *et al.*, 2014; Huang *et al.*, 2019), which suggests that *ERF019* may be involved in flg22-triggered immunity. To test this hypothesis, we used 3,3'-diaminobenzidine-tetrahydrochloride (DAB) staining to detect hydrogen peroxide accumulation in Col-0, the *erf019* mutant, and *ERF019*-OE lines on flg22 treatment. Although no obvious difference was observed between the *erf019* mutant and Col-0, our results revealed that the flg22-induced accumulation of hydrogen peroxide was impaired in *ERF019*-OE lines (OE71, OE72, and OE74) (Figure 6a). We also monitored the flg22-induced accumulation of ROS in *erf019* mutants and *ERF019*-OE lines. The results of this analysis were similar to those of DAB staining: ROS accumulation was impaired in *ERF019*-OE lines but not in the *erf019* mutants (Figure 6a). Moreover, we also found that the activation of MAPK3, MAPK4, and MAPK6, on flg22 treatment, was much stronger in both 267-31 and CRISPR/Cas9 mutants (ko13-14, ko26-14, and ko58-7) than in Col-0 (Figure 6c,d), and the PAMP-triggered MAPK activation was compromised in *ERF019*-OE lines




**FIGURE 5** Defence marker gene expression in the *erf019* mutants of *Arabidopsis thaliana*. Transcript levels of defence-related marker genes in the *erf019* mutants (267-31, ko13-4, and ko26-14) and Col-0 were evaluated by quantitative reverse transcription PCR (RT-qPCR) at different times postinoculation. RT-qPCR data are presented as relative transcript level for genes: *ICS1* and *PAL1*, two marker genes for salicylic acid (SA) biosynthesis; *PR1*, a marker for the SA signalling pathway; *LOX2*, a marker for jasmonic acid (JA) biosynthesis; *PDF1.2* and *VSP2*, involved in the JA signalling pathway; *FRK1*, a marker gene for the PAMP-triggered immunity (PTI) pathway; *ACS2* and *ACS6*, two marker genes for ethylene (ET) biosynthesis; and *ERF6* and *EIN2*, involved in the ET signalling pathway. *AtUBC9* was used as the internal control and transcript levels relative to Col-0 plants are displayed. Bars represent SE from three biological replicates and asterisks indicate statistical significance based on a two-tailed t test (\* $p < .05$ ; \*\* $p < .01$ ). 0 h, uninfected leaves

(OE71 and OE72) (Figure 6e). These results suggest that *ERF019* negatively regulates PAMP-triggered immunity.

To further demonstrate whether *ERF019* negatively regulates PTI triggered by *Phytophthora* elicitors, we tested whether it could inhibit INF1-induced necrosis. We first transiently overexpressed the *ERF019* protein in *N. benthamiana* leaves using *A. tumefaciens*-mediated transformation and found that *ERF019* did not induce necrosis after monitoring for up to 7 days postinfiltration (Figure S5). Next, we co-infiltrated mixtures of *A. tumefaciens* cultures carrying constructs of elicitors and either *ERF019* or FLAG-GFP into 5-week-old

*N. benthamiana* leaves. An HR in *N. benthamiana* leaves was observed 4 days postinfiltration, and the responses were classified into three categories according to the degree of response: no cell death, partial cell death, and full cell death (Figure 7a). Interestingly, *ERF019* expression significantly suppressed the HR induced by INF1 compared with the control FLAG-GFP (Figure 7b,c). We also co-infiltrated *ERF019* with the proapoptotic protein elicitor Bax and found that *ERF019* did not significantly inhibit Bax-induced cell death (Figure 7b,c). These results suggest that *ERF019* specifically suppresses *Phytophthora* elicitor INF1-triggered cell death in *N. benthamiana*. Taken together,



**FIGURE 6** Flg22-triggered immunity in *erf019* mutants and *ERF019* overexpression lines. (a) Detection of hydrogen peroxide by 3,3'-diaminobenzidine-tetrahydrochloride (DAB) staining. Leaves from the *erf019* mutant (267-31) and *ERF019* overexpression lines (OE71, OE72, and OE74) were infiltrated with 1  $\mu$ M flg22; hydrogen peroxide was detected 24 hr later. (b) The reactive oxygen species (ROS) burst in the 267-31 mutant and *ERF019* overexpression lines after treatment with 1  $\mu$ M flg22. Total relative luminescent units (RLUs) were detected over 30 min using leaf discs of 4-week-old plants. (c)–(e) Immunoblotting of phosphorylated MAP kinase on flg22 treatment in *erf019* mutants (*ko13-4*, *ko26-14*, *ko58-7*, and 267-31), *ERF019*-OE lines (OE71 and OE72), and wild-type Col-0. Samples were collected at 0, 10, and 60 min after flg22 treatment, total protein was extracted and analysed by immunoblots using antibodies against phospho-p44/42 MAPK and actin. Similar results were obtained in two independent experiments


we demonstrated that *ERF019* negatively regulates plant resistance by inhibiting PTI.

### 3 | DISCUSSION

The plant diseases caused by *Phytophthora* spp. pose a great threat to agriculture, highlighting the importance of studies on the

mechanisms of plant resistance. Previous studies based on forward genetics or map-based cloning technology have characterized dozens of R genes, which are commonly used in crop resistance breeding. However, R gene-mediated resistance, also described as effector recognition-based resistance, has been frequently overcome by new pathogen races. In contrast, disabling plant disease susceptibility genes (negative regulators of plant resistance) may provide a novel way to achieve durable and broad-spectrum resistance. For example,





**FIGURE 7** ERF019 suppresses INF1-triggered cell death in *Nicotiana benthamiana*. (a) The degree of elicitor-induced programmed cell death was categorized into three classes: no cell death (cell death area accounts for less than 30% of the injected area), partial cell death (cell death area accounts for 30%–80% of the injected area), and full cell death (cell death area accounts for more than 80% of the injected area). (b) Expression of ERF019 suppressed necrosis triggered by *Phytophthora infestans* PAMP elicitor INF1 but not Bax. *Agrobacterium tumefaciens* GV3101 cells carrying 35S::FLAG-GFP or 35S::ERF019 were mixed with cells carrying elicitor constructs and were co-infiltrated into 5-week-old *N. benthamiana* leaves. Phenotypic changes were monitored at 4 days postinfiltration. (c) Effect of ERF019 expression on necrosis triggered by INF1 and Bax. At least 30 infiltration sites were examined. Asterisks indicate statistical significance based on a two-tailed *t* test (\*\**p* < .01). Similar results were obtained in at least two independent experiments

loss-of-function of the *MLO* gene confers broad-spectrum resistance to the powdery mildew fungus (Jorgensen, 1992; Büschges *et al.*, 1997; Piffanelli *et al.*, 2002). The *mlo* allele has been introduced into European spring barley cultivars, and this has provided robust resistance for nearly four decades (Jorgensen, 1992; Lyngkjær and Carver, 2000), suggesting the great potential of disabling negative regulators in improving crop disease resistance.

Here, we showed the successful use of the compatible system between *P. parasitica* and *A. thaliana* to identify negative regulators. We identified the T-DNA insertion mutant 267-31, which was demonstrated to be resistant to *P. parasitica* without obvious inhibition of growth (Figure 1). Multiple *erf019* frameshift mutants generated by CRISPR/Cas9 technology consistently showed enhanced resistance to *P. parasitica*, while ERF019 overexpression lines were more susceptible (Figures 1 and 2). These results indicate that ERF019 negatively regulates plant resistance to *P. parasitica*.

The ERF family is a large family of transcription factors in plants, with up to 122 members in *Arabidopsis* and 139 members in rice (Nakano *et al.*, 2006). ERFs are involved in diverse developmental processes and various responses to environmental stimuli, such as pathogen attack, drought, salt, wounding, UV irradiation, and

extreme temperature (Tsutsui *et al.*, 2009; Liu *et al.*, 2012; Licausi *et al.*, 2013; Maruyama *et al.*, 2013). ERF019 was classified into phylogenetic Group II of the *Arabidopsis* ERF family (Nakano *et al.*, 2006). There are 15 members in this group, which are further classified into three subgroups: IIa, IIb, and IIc. Most genes in subgroups IIa and IIb have been shown to play crucial roles in biotic and abiotic stress responses. For example, overexpression of *DEAR1*, a subgroup IIa gene, rendered *Arabidopsis* more resistant to *Pseudomonas syringae* infection and less tolerant to freezing (Tsutsui *et al.*, 2009). Furthermore, all six genes of subgroup IIa induce cell death in tobacco (Ogata *et al.*, 2013). ERF15, a subgroup IIb gene, has been reported to be a negative regulator of salt and drought tolerance (Lee *et al.*, 2015). Here, we found that the subgroup IIc gene ERF019 negatively regulates plant resistance to *P. parasitica*. It has also been reported that overexpression of ERF019 delays plant growth and senescence, enhances drought resistance, and increases plant susceptibility to *Botrytis cinerea* and *P. syringae* (Scarpeci *et al.*, 2017; Huang *et al.*, 2019). These observations suggest that ERF family subgroup II members may play important roles in plant resistance and abiotic stress, and functional analysis of their orthologous genes in crops will provide potential gene resources for breeding for disease resistance.

Plant PTI responses, including ROS burst, callose deposition, MAP kinase activity, and defence gene induction, are critical for plants to repel pathogen attacks (Bigeard *et al.*, 2015). We show here that the induction of the PTI-related marker gene *FRK1* in the 267-31 mutant on *P. parasitica* infection was stronger than that in the wild type Col-0 (Figure 5). Meanwhile, flg22-induced activation of the MPK3, MPK4, and MPK6 was enhanced in both the 267-31 mutant and *erf019* knockout lines when compared with Col-0, and was compromised in *ERF019*-OE lines (Figure 6). The flg22-induced ROS burst and hydrogen peroxide accumulation were also significantly suppressed in the leaves of *ERF019*-OE plants (Figure 6), which is consistent with the previous report that flg22-induced callose deposition was significantly impaired in *ERF019*-OE plants (Huang *et al.*, 2019). Furthermore, we showed that transient overexpression of *ERF019* in *N. benthamiana* suppresses *Phytophthora* PAMP elicitor *INF1*-activated cell death (Figure 7). These results demonstrate the critical role of *ERF019* in negatively regulating PTI responses.

Plant cell death plays a central role in interactions with hemibiotrophic pathogens, such as *Phytophthora* species, considering that these pathogens initially develop haustoria to acquire nutrients from living host cells and then subsequently switch to a necrotrophic lifestyle, resulting in the death of the host plant (Lamour *et al.*, 2012). *INF1*-like proteins are a family of secreted elicitors, which exist widely in *Phytophthora*. It is reasonable to hypothesize that *ERF019* may negatively regulate plant resistance by suppressing cell death, which facilitates the growth of *P. parasitica* during plant infection. Meanwhile, *ERF019* cannot inhibit Bax-induced cell death, like *P. sojae* effector Avh238 that was reported to inhibit *INF1*- but not Bax-induced cell death (Wang *et al.*, 2011a). Bax is a proapoptotic member and can translocate into the mitochondrial membrane and trigger the apoptotic process, some features of which resemble plant programmed cell death (Ihara-Ohori *et al.*, 2007). However, compared to Bax-induced cell death, the recognition of *INF1* and the downstream pathway of *INF1*-triggered cell death may possess some unique features, some of which may be regulated by *ERF019*. Because the overexpression of *ERF019* leads to attenuation of PTI responses, and the silencing of *N. benthamiana* receptor-like kinase gene *SERK3*, which encodes a homolog of *Arabidopsis* BAK1 and plays a key role in PTI by suppressing *INF1*-induced cell death (Chaparro-Garcia *et al.*, 2011), it is likely that *ERF019* suppresses *INF1*-activated cell death by interfering with the PTI signalling pathway.


Interestingly, the negative regulator *ERF019* was induced during *P. parasitica* infection. Moreover, *ERF019* was also highly induced by flg22 (Huang *et al.*, 2019). These observations suggested that *ERF019* can be induced during PTI, which in turn inhibits PTI. Previous researchers showed that plant recognition of PAMPs induces both positive and negative PTI signalling pathways. For example, the PAMP-induced MEKK1, MEKK1/2, and MPK4 signalling cascades negatively mediate plant defence responses (Ichimura *et al.*, 2006; Mészáros *et al.*, 2006; Suarez-Rodriguez *et al.*, 2007; Gao *et al.*, 2008; Qiu *et al.*, 2008; Pitzschke *et al.*, 2009). Thus, *ERF019* may

be involved in a negative feedback loop that balances growth and resistance.

Our results also showed that the SA biosynthesis-related gene *ICS1*, SA signalling pathway marker gene *PR1*, JA signalling marker gene *VSP2* and *PDF1.2* as well as JA biosynthesis-related gene *LOX2* were up-regulated in the *erf019* mutant, indicating that the SA and JA signalling pathways are coupled through *ERF019*. It has been reported that SA is responsible for plant defence against biotrophs, whereas JA or ET is responsible for defence against necrotrophs (Bostock, 2005). However, both the SA and JA signalling pathways have been shown to contribute to basal resistance against *P. parasitica* (Attard *et al.*, 2010). Interference with SA, JA, or ET signalling in the *eds1*, *eds5*, *pad4*, *sid2*, *ein2*, *etr1*, and *jar1* mutants and *NahG* transgenic plants enhanced plant susceptibility to *P. parasitica* (Attard *et al.*, 2010). Moreover, the *Arabidopsis thaliana* Resistant to *Phytophthora* 5 gene (*AtrTP5*), which encodes a WD40 repeat domain-containing protein, has been reported to negatively regulate plant resistance to *P. parasitica* by interfering with the JA and SA signalling pathways (Li *et al.*, 2020). In addition, *ERF019* was shown to be induced by OPDA, a cyclopentenone precursor of JA (Taki *et al.*, 2005). The function of *ERF019* may be repressed by NINJA (the transcriptional co-repressor Novel INTERactor of JAZ), a negative regulator of JA signalling, through protein-protein interaction (Huang *et al.*, 2019). These results suggest a potential role of *ERF019* in the JA signalling pathway, which is subjected to complex positive and negative regulation and is coupled with the SA signalling pathway.

Loss of function of a negative regulator of plant resistance may constitutively activate defence responses and reduce plant fitness (Tian *et al.*, 2003; Denancé *et al.*, 2013; Huot *et al.*, 2014). For example, loss of function of the *MPK4* gene results in a dwarf phenotype, which is accompanied by elevated SA levels and constitutive expression of pathogenesis-related genes (Petersen *et al.*, 2000). The *mek1* mutant and *mkk1 mkk2* double mutant also display constitutive defence responses and reduced plant growth (Gao *et al.*, 2008). However, *erf019* plants exhibit resistance to *P. parasitica* without altered plant growth. Consistent with this phenotype, RT-qPCR results showed that the expression of *ERF019* was low in adult rosette leaves under normal conditions (Figure 3), suggesting that *ERF019* may not be necessary for plant growth. Furthermore, the expression of defence-related marker genes was just slightly up-regulated in the *erf019* mutant in the absence of pathogen infection (Figure 5). In addition, after flg22 treatment, activation of MAP kinase was stronger in the *erf019* mutants than in the wild-type Col-0 and attenuated in the *ERF019*-OE plants when compared to the wild-type Col-0 after 10 min of treatment (Figure 6c,d,e). These observations show that PTI responses seem to be amplified in the *erf019* mutant without strong constitutive induction of the expression of pathogenesis-related genes, thus increasing resistance without influencing growth.

*ERF019* contains a conserved AP2/ERF DNA-binding domain at the N-terminus. There was no conserved motif identified at the C-terminus of *ERF019* based on multiple sequence alignment



**FIGURE 8** Proposed model for the role of *ERF019* in plant immunity. The PAMP-triggered immunity (PTI) and multiple defence pathways, including the salicylic acid (SA) and jasmonic acid (JA) signal pathways, are induced on infection or flg22-treatment. *ERF019* is also induced on infection by *Phytophthora parasitica* but suppresses PTI as well as SA and JA signal pathways, suggesting it plays an important role in the negative feedback loop that balances growth and resistance

analyses of ERF family members (Nakano *et al.*, 2006). The potentially accumulated proteins in the CRISPR/Cas9-edited *erf019* mutant lines, which are predicted to be truncated due to the frameshift mutations from the codon for the 109th amino acid, lost ability to negatively regulate resistance, indicating a key role of the C-terminus of *ERF019* in immune function. The regions outside the DNA-binding domain in ERF proteins are generally involved in protein modification and protein-protein interactions, and are important for their nuclear localization and transcriptional activities (Nakano *et al.*, 2006; Licausi *et al.*, 2013). For example, the C-terminal activation domain, but not the N-terminal DNA binding domain of ERF protein TINY, interacts with and antagonizes BRASSINOSTEROID INSENSITIVE1-ETHYL METHANESULFONATE SUPPRESSOR1 (BES1) in the regulation of drought response (Xie *et al.*, 2019). The transcriptional repressor Novel INteractor of JAZ (NINJA) interacts with *ERF019* and represses its function (Huang *et al.*, 2019). We speculate that the C-terminus of *ERF019* mediates interaction by other unknown protein factors to regulate plant immunity. Future efforts to identify the *ERF019*-interacting proteins will be useful in understanding the underlying mechanisms of *ERF019* in regulating plant immunity.

Based on our study, we propose that *ERF019* plays an important role in the negative feedback loop that balances growth and resistance on pathogen infection by suppressing PTI and SA/JA defence responses (Figure 8). Further identification of target genes regulated by *ERF019* will provide insights into the mechanisms of the negative feedback loop. Identification of loss-of-function alleles of *ERF019* and its homologs in crops is a potential strategy for breeding crops with durable resistance.

## 4 | EXPERIMENTAL PROCEDURES

### 4.1 | Plant materials and growth conditions

The *Arabidopsis* T-DNA insertion lines were generated (Zhang *et al.*, 2005) and kindly provided by Dr Jianru Zuo. *Arabidopsis* ecotype Col-0 and T-DNA insertion mutants used in this study were grown at 23°C with 14 hr of light per 24 hr. *N. benthamiana* plants were also grown at 23°C with 14 hr of light per 24 hr.

### 4.2 | Pathogen growth and infection assays

*P. parasitica* strain Pp016 was originally isolated from diseased tobacco plants in Queensland, Australia (Wang *et al.*, 2011b; Zhang *et al.*, 2011), and 1121 is a transformant of *P. parasitica* Pp016 stably expressing ER-rendered GFP under the control of the constitutive *Hsp70* promoter of *B. lactucae*. The transformant 1121 remained pathogenic on *A. thaliana* and tobacco plants, similar to the wild-type strain Pp016 (Zhang *et al.*, 2011). The GFP in *P. parasitica* 1121 is constitutively and widely expressed in the cytoplasm, allowing easy monitoring during colonization of host plants. *P. parasitica* culture conditions, zoospore production, and the detached leaf inoculation assays were performed as described (Wang *et al.*, 2011b). For the pathogenicity assay, the detached fully expanded apical leaves from approximately 4-week-old *A. thaliana* plants were wounded with a toothpick and the zoospore suspensions were adjusted to a concentration of 200 zoospores/ $\mu$ l using microscopy and applied as droplets at the wounding sites to ensure infection. To observe the lesion size more clearly, trypan blue was used to stain the death plant cells in lesions of inoculated leaves as described (Li *et al.*, 2019). *A. thaliana* leaves infected with the *P. parasitica* transformant 1121 were observed with the OLYMPUS BX51 fluorescence microscope (with excitation at 450–480 nm and emission at 515 nm) to detect *P. parasitica* hyphae (green fluorescence) at 2 dpi. For pathogen biomass, three biological replicates were performed with at least eight leaves per replicate. Primers used for pathogen biomass were listed in Table S1. Disease severity was evaluated based on the lesion sizes on detached leaves and the extent of pathogen colonization.

### 4.3 | TAIL-PCR and RT-qPCR assays

TAIL-PCR was performed as described (Liu *et al.*, 1995). For RT-qPCR assays, three biological replicates were used. Total RNA from the whole leaves before and after *P. parasitica* inoculation was extracted using TRIzol reagent (Invitrogen). One microgram of total RNA was used to perform reverse transcription using the PrimeScript RT Reagent Kit with gDNA Eraser (TAKARA). For real-time qPCR analysis, 0.5  $\mu$ l of the first-strand cDNA reaction products was used as template in a reaction with Ultra SYBR Mixture (CW BIO) under the following conditions: 95°C for 10 min, and 40 cycles of 95°C for 15 s and 60°C for 30 s. The

fold changes in target gene expression were normalized using *UBC9* as the internal control. Primers used for RT-qPCR are listed in Table S1.

#### 4.4 | CRISPR/Cas9-based knockouts and overexpression of *ERF019* in *Arabidopsis*

For the 35S::*ERF019* construct, full-length *ERF019* was directionally cloned into pKANNIBAL (Wesley *et al.*, 2001) and then subcloned into the binary vector pART27 (Gleave, 1992). For the CRISPR/Cas9-based knockouts, two 19-bp sgRNA oligonucleotides targeting the exon of *ERF019* were inserted in the psgR-Cas9 vector to create deletion mutants as previously described (Feng *et al.*, 2013, 2014). Annealed 19-bp sgRNA oligomers were inserted into the *BbsI* site of the psgR-Cas9 vector. Based on the psgR-Cas9 vector, the second pATU6-sgR cassette was amplified by PCR after the insertion of target oligomers and ligated into the *KpnI/EcoRI* site of the above psgR-Cas9 vector. The cassette was then transferred into the binary vector pCXSN. The generated binary vectors were transformed into *A. tumefaciens* GV3101. *A. tumefaciens* cells carrying 35S::*ERF019* and sgRNAs/Cas9 constructs were transformed into wild-type Col-0 via the floral-dip method (Zhang *et al.*, 2006). 35S::*ERF019* transformants were screened on 1/2 Murashige and Skoog (MS) agar plates containing 50 µg/ml kanamycin. sgRNAs/Cas9 transformants were screened on 1/2 MS agar plates containing 50 µg/ml hygromycin.

#### 4.5 | *A. tumefaciens* infiltration assays

*A. tumefaciens* GV3101 containing constructs was cultured at 28°C and 200 rpm for approximately 24 hr in Luria Bertani (LB) medium with appropriate antibiotics. The *A. tumefaciens* cells were collected by centrifugation, resuspended in infiltration buffer (10 mM MES, 10 mM MgCl<sub>2</sub>, 0.2 mM acetosyringone, pH 5.6), and adjusted to the appropriate OD<sub>600</sub> before being infiltrated into 5-week-old *N. benthamiana* leaves (the OD<sub>600</sub> was generally 0.3 for confocal subcellular localization assays and 0.15–0.4 for infection assays). For infection assays, *A. tumefaciens* GV3101 cells carrying constructs were suspended at a concentration of OD<sub>600</sub> = 0.4, and then inoculated with *P. parasitica* 2 days after infiltration. For co-expression assays, *A. tumefaciens* GV3101 cells carrying constructs were suspended at an appropriate OD<sub>600</sub> and mixed before infiltration (for INF1 OD<sub>600</sub> = 0.15, for Bax OD<sub>600</sub> = 0.3, for FLAG-GFP and *ERF019* OD<sub>600</sub> = 0.4). Symptom development was monitored visually from 2 to 6 days after infiltration depending on the cell death activator.

#### 4.6 | Subcellular localization assays

*A. tumefaciens* GV3101 cell cultures carrying constructs expressing FLAG-GFP, *ERF019*-GFP-NES, *ERF019*-GFP-nes, *ERF019*-GFP or nucleus marker H2B-mCherry were collected by centrifugation and resuspended in infiltration buffer (10 mM MES, 10 mM MgCl<sub>2</sub>, 0.2 mM

acetosyringone, [pH 5.6]) at OD<sub>600</sub> = 0.4 and then the cells carrying GFP constructs were co-infiltrated with those expressing the nucleus marker H2B-mCherry into 5-week-old *N. benthamiana* leaves. Three days after infiltration, the *N. benthamiana* leaves were visually inspected under an Olympus FV3000 confocal microscope with excitation wavelengths of 488 nm for GFP and 587 nm for mCherry.

#### 4.7 | Detection of hydrogen peroxide

Hydrogen peroxide was detected in *Arabidopsis* rosette leaves infiltrated with 1 µM flg22 as described previously (Daudi and O'Brien, 2012).

#### 4.8 | Oxidative burst measurements

ROS was measured in 30-day-old *Arabidopsis* seedlings. In brief, the leaf disks were cut from 30-day-old mature leaves with a sharp 5-mm puncher and were floated in sterile ultrapure water in culture dishes overnight. The next day, the leaf disks were transferred into 96-well plates and 100 µl Luminol Enhancer (CW BIO, CW0049M) and 100 µl of 20 µg/ml horseradish peroxidase (Aladdin) were added into each cell. Then, 5 µl of 41 µM flg22 was immediately added into each cell to a final concentration of 1 µM. Luminescence was measured using a TECAN Infinite M200 PRO (TECAN); 40 cycles (1 min per cycle) were used for the measurement.

#### 4.9 | MAPK activity assays

*Arabidopsis* seedlings were grown on vertical 1/2 MS plates at 23°C with 14 hr of light per 24 hr for 12 days and then were transferred into liquid MS and incubated overnight with minimum rotation speed (40 rpm). The next day, the seedlings were treated with 1 µM flg22 and frozen in liquid nitrogen. Total proteins were extracted with glycerol-Tris-EDTA-NaCl buffer (10% glycerol, 25 mM Tris-HCl [pH 7.5], 1 mM EDTA, 150 mM NaCl, 0.1% Tween 20, 0.1% NP-40, 2% [vol/vol] PVPP, 0.1 mM DTT, 1 × inhibitor cocktail, 1 × phosphatase inhibitor cocktail 2 and 1 × phosphatase inhibitor cocktail 3). The protein concentration was measured using the Super-Bradford Protein Assay Kit (CW BIO, CW0013S). Equal amounts of total protein were loaded on a 10% SDS-PAGE gel. Anti-phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (D13.14.4E) XP rabbit mAb antibody (Cell Signaling Technology) was used to detect the phosphorylation state of MPK3, MPK4, and MPK6.

#### ACKNOWLEDGMENTS

We would like to thank Professor Jim Peacock and Professor Liz Dennis (CSIRO Agriculture and Food, Canberra, Australia), Professor Katrina M. Ramonell (University of Alabama, USA) and Professor Gary Loake (University of Edinburgh, UK) for their helpful suggestions on the manuscript, and Northwest A&F University Life Science Research Core Services for providing advanced facilities. This work was supported by the National Natural Science Foundation of China

(31125022 and 31930094), the China Agriculture Research System (CARS-09), and the Programme of Introducing Talents of Innovative Discipline to Universities (project 111) from the State Administration of Foreign Experts Affairs (#B18042).

#### AUTHOR CONTRIBUTIONS

W.S., Y.M., and W.L. designed the experiments. W.L., F.D., J.J., X.C., J.L., Q.W., T.L., and Y.M. performed the experiments. W.L., J.J., X.C., Y.M., and W.S. analysed the data. W.L., Y.M., and W.S. wrote the manuscript with contributions from all authors.

#### DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

#### ORCID

Weixing Shan  <https://orcid.org/0000-0001-7286-4041>

#### REFERENCES

- Albrecht, C., Boutrot, F., Segonzac, C., Schwessinger, B., Gimenez-Ibanez, S., Chinchilla, D. *et al.* (2012) Brassinosteroids inhibit pathogen-associated molecular pattern-triggered immune signaling independent of the receptor kinase BAK1. *Proceedings of the National Academy of Sciences of the United States of America*, 109, 303–308.
- Alonso, J.M., Hirayama, T., Roman, G., Nourizadeh, S. and Ecker, J.R. (1999) EIN2, a bifunctional transducer of ethylene and stress responses in *Arabidopsis*. *Science*, 284, 2148–2152.
- Attard, A., Gourgues, M., Callemeyn-Torre, N. and Keller, H. (2010) The immediate activation of defence responses in *Arabidopsis* roots is not sufficient to prevent *Phytophthora parasitica* infection. *New Phytologist*, 187, 449–460.
- Bigeard, J., Colcombet, J. and Hirt, H. (2015) Signaling mechanisms in pattern-triggered immunity (PTI). *Molecular Plant Pathology*, 8, 521–539.
- Boller, T. and Felix, G. (2009) A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annual Review of Plant Biology*, 60, 379–406.
- Bostock, R.M. (2005) Signal crosstalk and induced resistance: straddling the line between cost and benefit. *Annual Review of Phytopathology*, 43, 545–580.
- Boutrot, F. and Zipfel, C. (2017) Function, discovery, and exploitation of plant pattern recognition receptors for broad-spectrum disease resistance. *Annual Review of Phytopathology*, 55, 257–286.
- Büschges, R., Hollricher, K., Panstruga, R., Simons, G., Wolter, M., Frijters, A. *et al.* (1997) The barley *Mlo* gene: a novel control element of plant pathogen resistance. *Cell*, 88, 695–705.
- Chaparro-Garcia, A., Wilkinson, R.C., Gimenez-Ibanez, S., Findlay, K., Coffey, M.D., Zipfel, C. *et al.* (2011) The receptor-like kinase SERK3/BAK1 is required for basal resistance against the late blight pathogen *Phytophthora infestans* in *Nicotiana benthamiana*. *PLoS ONE*, 6, e16608.
- Chen, Z., Agnew, J.L., Cohen, J.D., He, P., Shan, L., Sheen, J. *et al.* (2007) *Pseudomonas syringae* type III effector AvrRpt2 alters *Arabidopsis thaliana* auxin physiology. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 20131–20136.
- Daudi, A. and O'Brien, J.A. (2012) Detection of hydrogen peroxide by DAB staining in *Arabidopsis* leaves. *Bio-Protocols*, 2, e263.
- Denancé, N., Sánchez-Vallet, A., Goffner, D. and Molina, A. (2013) Disease resistance or growth: the role of plant hormones in balancing immune responses and fitness costs. *Frontiers in Plant Science*, 4, 155.
- Deng, Y., Zhai, K., Xie, Z., Yang, D., Zhu, X., Liu, J. *et al.* (2017) Epigenetic regulation of antagonistic receptors confers rice blast resistance with yield balance. *Science*, 355, 962–965.
- Feng, Z., Zhang, B., Ding, W., Liu, X., Yang, D.L., Wei, P. *et al.* (2013) Efficient genome editing in plants using a CRISPR/Cas system. *Cell Research*, 23, 1229–1232.
- Feng, Z., Mao, Y., Xu, N., Zhang, B., Wei, P., Yang, D.L. *et al.* (2014) Multigeneration analysis reveals the inheritance, specificity, and patterns of CRISPR/Cas-induced gene modifications in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America*, 111, 4632–4637.
- Gao, M., Liu, J., Bi, D., Zhang, Z., Cheng, F., Chen, S. *et al.* (2008) MEKK1, MKK1/MKK2 and MPK4 function together in a mitogen-activated protein kinase cascade to regulate innate immunity in plants. *Cell Research*, 18, 1190–1198.
- Gleave, A.P. (1992) A versatile binary vector system with a T-DNA organisational structure conducive to efficient integration of cloned DNA into the plant genome. *Plant Molecular Biology*, 20, 1203–1207.
- Grünwald, N.J., Garbelotto, M., Goss, E.M., Heungens, K. and Prospero, S. (2012) Emergence of the sudden oak death pathogen *Phytophthora ramorum*. *Trends in Microbiology*, 20, 131–138.
- Haverkort, A.J., Boonekamp, P.M., Hutten, R., Jacobsen, E., Lotz, L.A.P., Kessel, G.J.T. *et al.* (2008) Societal costs of late blight in potato and prospects of durable resistance through cisgenic modification. *Potato Research*, 51, 47–57.
- Huang, P., Zhang, J., Jiang, B., Chan, C., Yu, J., Lu, Y. *et al.* (2019) NINJA-associated ERF19 negatively regulates *Arabidopsis* pattern-triggered immunity. *Journal of Experimental Botany*, 70, 1033–1047.
- Huot, B., Yao, J., Montgomery, B.L. and He, S.Y. (2014) Growth-defense tradeoffs in plants: a balancing act to optimize fitness. *Molecular Plant Pathology*, 7, 1267–1287.
- Ichimura, K., Casais, C., Peck, S.C., Shinozaki, K. and Shirasu, K. (2006) MEKK1 is required for MPK4 activation and regulates tissue-specific and temperature-dependent cell death in *Arabidopsis*. *Journal of Biological Chemistry*, 281, 36969–36976.
- Ihara-Ohori, Y., Nagano, M., Muto, S., Uchimiya, H. and Kawai-Yamada, M. (2007) Cell death suppressor *Arabidopsis* bax inhibitor-1 is associated with calmodulin binding and ion homeostasis. *Plant Physiology*, 143, 650–660.
- Jaillais, Y. and Vert, G. (2012) Brassinosteroids, gibberellins and light-mediated signalling are the three-way controls of plant sprouting. *Nature Cell Biology*, 14, 788–790.
- Jones, J.D. and Dangl, J.L. (2006) The plant immune system. *Nature*, 444, 323–329.
- Jorgensen, J.H. (1992) Discovery, characterization and exploitation of *Mlo* powdery mildew resistance in barley. *Euphytica*, 63, 141–152.
- Kamoun, S., Furzer, O., Jones, J.D.G., Judelson, H.S., Ali, G.S., Dhalio, R.J.D. *et al.* (2015) The top 10 oomycete pathogens in molecular plant pathology. *Molecular Plant Pathology*, 16, 413–434.
- Lamour, K.H., Stam, R., Jupe, J. and Huitema, E. (2012) The oomycete broad-host-range pathogen *Phytophthora capsici*. *Molecular Plant Pathology*, 13, 329–337.
- Lee, S., Lee, S. and Kim, S.Y. (2015) AtERF15 is a positive regulator of ABA response. *Plant Cell Reports*, 34, 71–81.
- Li, T., Wang, Q., Feng, R., Li, L., Ding, L., Fan, G. *et al.* (2019) Negative regulators of plant immunity derived from cinnamyl alcohol dehydrogenases are targeted by multiple *Phytophthora* Avr3a-like effectors. *New Phytologist*, <https://doi.org/10.1111/nph.16139>
- Li, W., Zhao, D., Dong, J., Kong, X., Zhang, Q., Li, T. *et al.* (2020) AtRTP5 negatively regulates plant resistance to *Phytophthora* pathogens by modulating the biosynthesis of endogenous jasmonic acid and salicylic acid. *Molecular Plant Pathology*, 21, 95–108.
- Licausi, F., Ohme-Takagi, M. and Perata, P. (2013) APETALA2/Ethylene Responsive Factor (AP2/ERF) transcription factors: mediators of

- stress responses and developmental programs. *New Phytologist*, 199, 639–649.
- Liu, D., Chen, X., Liu, J., Ye, J. and Guo, Z. (2012) The rice ERF transcription factor OsERF922 negatively regulates resistance to *Magnaporthe oryzae* and salt tolerance. *Journal of Experimental Botany*, 63, 3899–3911.
- Liu, Y.G., Mitsukawa, N., Oosumi, T. and Whittier, R.F. (1995) Efficient isolation and mapping of *Arabidopsis thaliana* T-DNA insert junctions by thermal asymmetric interlaced PCR. *The Plant Journal*, 8, 457–463.
- Lyngkjær, M.F. and Carver, T.L. (2000) Conditioning of cellular defence responses to powdery mildew in cereal leaves by prior attack. *Molecular Plant Pathology*, 1, 41–49.
- Maruyama, Y., Yamoto, N., Suzuki, Y., Chiba, Y., Yamazaki, K., Sato, T. et al. (2013) The *Arabidopsis* transcriptional repressor ERF9 participates in resistance against necrotrophic fungi. *Plant Science*, 213, 79–87.
- Meng, Y., Zhang, Q., Ding, W. and Shan, W. (2014) *Phytophthora parasitica*: a model oomycete plant pathogen. *Mycology*, 5, 43–51.
- Mészáros, T., Helfer, A., Hatzimasoura, E., Magyar, Z., Serazetdinova, L., Rios, G. et al. (2006) The *Arabidopsis* MAP kinase kinase MKK1 participates in defence responses to the bacterial elicitor flagellin. *The Plant Journal*, 48, 485–498.
- Moffat, C.S., Ingle, R.A., Wathugala, D.L., Saunders, N.J., Knight, H. and Knight, M.R. (2012) ERF5 and ERF6 play redundant roles as positive regulators of JA/ET-mediated defense against *Botrytis cinerea* in *Arabidopsis*. *PLoS ONE*, 7, e35995.
- Murphy, F., He, Q., Armstrong, M., Giuliani, L.M., Boevink, P.C., Zhang, W. et al. (2018) The potato MAP3K StVIK1s is required for the *Phytophthora infestans* RXLR effector Pi17316 to promote disease. *Plant Physiology*, 177, 398–410.
- Naito, K., Taguchi, F., Suzuki, T., Inagaki, Y., Toyoda, K., Shiraishi, T. et al. (2008) Amino acid sequence of bacterial microbe-associated molecular pattern flg22 is required for virulence. *Molecular Plant-Microbe Interactions*, 21, 1165–1174.
- Nakano, T., Suzuki, K., Fujimura, T. and Shinshi, H. (2006) Genome-wide analysis of the ERF gene family in *Arabidopsis* and rice. *Plant Physiology*, 140, 411–432.
- Ogata, T., Kida, Y., Tochigi, M. and Matsushita, Y. (2013) Analysis of the cell death-inducing ability of the ethylene response factors in group VIII of the AP2/ERF family. *Plant Science*, 209, 12–23.
- Panabières, F., Ali, G.S., Allagui, M.B., Dalio, R.J.D., Gudmestad, N.C., Kuhn, M.L. et al. (2016) *Phytophthora nicotianae* diseases worldwide: New knowledge of a long-recognised pathogen. *Phytopathologia Mediterranea*, 55, 20–40.
- Petersen, M., Brodersen, P., Naested, H., Andreasson, E., Lindhart, U., Johansen, B. et al. (2000) *Arabidopsis* MAP kinase 4 negatively regulates systemic acquired resistance. *Cell*, 103, 1111–1120.
- Pieterse, C.M.J., Leon-Reyes, A., Van der Ent, S. and Van Wees, S.C.M. (2009) Networking by small-molecule hormones in plant immunity. *Nature Chemical Biology*, 5, 308–316.
- Piffanelli, P., Zhou, F., Casais, C., Orme, J., Jarosch, B., Schaffrath, U. et al. (2002) The barley MLO modulator of defense and cell death is responsive to biotic and abiotic stress stimuli. *Plant Physiology*, 129, 1076–1085.
- Pitzschke, A., Schikora, A. and Hirt, H. (2009) MAPK cascade signaling networks in plant defence. *Current Opinion in Plant Biology*, 12, 421–426.
- Qiu, J., Fiil, B.K., Petersen, K., Nielsen, H.B., Botanga, C.J., Thorgrimsen, S. et al. (2008) *Arabidopsis* MAP kinase 4 regulates gene expression through transcription factor release in the nucleus. *EMBO Journal*, 27, 2214–2221.
- Sano, S., Aoyama, M., Nakai, K., Shimotani, K., Yamasaki, K., Sato, M.H. et al. (2014) Light-dependent expression of flg22-induced defense genes in *Arabidopsis*. *Frontiers in Plant Science*, 5, 531.
- Sasaki, Y., Asamizu, E., Shibata, D., Nakamura, Y., Kaneko, T., Awai, K. et al. (2001) Monitoring of methyl jasmonate-responsive genes in *Arabidopsis* by cDNA macroarray: self-activation of jasmonic acid biosynthesis and crosstalk with other phytohormone signaling pathways. *DNA Research*, 8, 153–161.
- Scarpeci, T.E., Frea, V.S., Zanor, M.I. and Valle, E.M. (2017) Overexpression of AtERF019 delays plant growth and senescence, and improves drought tolerance in *Arabidopsis*. *Journal of Experimental Botany*, 68, 673–685.
- Shan, L., He, P., Li, J., Heese, A., Peck, S.C., Nürnberger, T. et al. (2008) Bacterial effectors target the common signaling partner BAK1 to disrupt multiple MAMP receptor-signaling complexes and impede plant immunity. *Cell Host & Microbe*, 4, 17–27.
- Suarez-Rodriguez, M.C., Adams-Phillips, L., Liu, Y., Wang, H., Su, S., Jester, P.J. et al. (2007) MEKK1 is required for flg22-induced MPK4 activation in *Arabidopsis* plants. *Plant Physiology*, 143, 661–669.
- Taki, N., Sasaki-Sekimoto, Y., Obayashi, T., Kikuta, A., Kobayashi, K., Ainai, T. et al. (2005) 12-oxo-phytodienoic acid triggers expression of a distinct set of genes and plays a role in wound-induced gene expression in *Arabidopsis*. *Plant Physiology*, 139, 1268–1283.
- Tian, D., Traw, M.B., Chen, J.Q., Kreitman, M. and Bergelson, J. (2003) Fitness costs of R-gene-mediated resistance in *Arabidopsis thaliana*. *Nature*, 423, 74–77.
- Tsutsui, T., Kato, W., Asada, Y., Sako, K., Sato, T., Sonoda, Y. et al. (2009) DEAR1, a transcriptional repressor of DREB protein that mediates plant defense and freezing stress responses in *Arabidopsis*. *Journal of Plant Research*, 122, 633–643.
- Tyler, B.M. (2002) Molecular basis of recognition between *Phytophthora* pathogens and their hosts. *Annual Review of Phytopathology*, 40, 137–167.
- Uknes, S., Mauch-Mani, B., Moyer, M., Potter, S., Williams, S., Dincher, S. et al. (1992) Acquired resistance in *Arabidopsis*. *The Plant Cell*, 4, 645–656.
- Van der Straeten, D., Rodrigues-Pousada, R.A., Villarreal, R., Hanley, S., Goodman, H.M. and Van Montagu, M. (1992) Cloning, genetic mapping, and expression analysis of an *Arabidopsis thaliana* gene that encodes 1-aminocyclopropane-1-carboxylate synthase. *Proceedings of the National Academy of Sciences of the United States of America*, 89, 9969–9973.
- Wang, Q., Han, C., Ferreira, A.O., Yu, X., Ye, W., Tripathy, S. et al. (2011a) Transcriptional programming and functional interactions within the *Phytophthora sojae* RXLR effector repertoire. *The Plant Cell*, 23, 2064–2086.
- Wang, Y., Meng, Y., Zhang, M., Tong, X., Wang, Q., Sun, Y. et al. (2011b) Infection of *Arabidopsis thaliana* by *Phytophthora parasitica* and identification of variation in host specificity. *Molecular Plant Pathology*, 12, 187–201.
- Wang, X., Boevink, P., McLellan, H., Armstrong, M., Bukharova, T., Qin, Z. et al. (2015) A host KH RNA-binding protein is a susceptibility factor targeted by an RXLR effector to promote late blight disease. *Molecular Plant Pathology*, 8, 1385–1395.
- Wesley, S.V., Helliwell, C.A., Smith, N.A., Wang, M.B., Rouse, D.T., Liu, Q. et al. (2001) Construct design for efficient, effective and high-throughput gene silencing in plants. *The Plant Journal*, 27, 581–590.
- Wildermuth, M.C., Dewdney, J., Wu, G. and Ausubel, F.M. (2001) Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature*, 414, 562–565.
- Xie, Z., Nolan, T., Jiang, H., Tang, B., Zhang, M., Li, Z. et al. (2019) The AP2/ERF transcription factor TINY modulates brassinosteroid-regulated plant growth and drought responses in *Arabidopsis*. *The Plant Cell*, 31, 1788–1806.
- Yamada, T. (1993) The role of auxin in plant-disease development. *Annual Review of Phytopathology*, 31, 253–273.

- Zhang, J., Xu, J., Kong, Y., Ji, Z., Wang, X., An, F. *et al.* (2005) Generation of chemical-inducible activation tagging T-DNA insertion lines of *Arabidopsis thaliana*. *Acta Genetica Sinica*, 32, 1082–1088.
- Zhang, X., Henriques, R., Lin, S., Niu, Q. and Chua, N. (2006) *Agrobacterium*-mediated transformation of *Arabidopsis thaliana* using the floral dip method. *Nature Protocols*, 1, 641–646.
- Zhang, M., Wang, Q., Xu, K., Meng, Y., Quan, J. and Shan, W. (2011) Production of dsRNA sequences in the host plant is not sufficient to initiate gene silencing in the colonizing oomycete pathogen *Phytophthora parasitica*. *PLoS ONE*, 6, e28114–e28210.
- Zipfel, C., Robatzek, S., Navarro, L., Oakeley, E.J., Jones, J.D.G., Felix, G. *et al.* (2004) Bacterial disease resistance in *Arabidopsis* through flagellin perception. *Nature*, 428, 764–767.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Lu W, Deng F, Jia J, et al. The *Arabidopsis thaliana* gene *AtERF019* negatively regulates plant resistance to *Phytophthora parasitica* by suppressing PAMP-triggered immunity. *Molecular Plant Pathology*. 2020;21:1179–1193. <https://doi.org/10.1111/mpp.12971>